



DISINFECTING PROTOCOL

Always remove animals, food and water dishes, bedding and toys from the area before implementing the cleaning protocol.

DIRECTIONS FOR USE:

1. Remove cap from bottle.
2. Squeeze bottle until the desired level.
3. Pour concentrate into the container.
4. Add water.



*DILUTION RATIO & EFFICACY

1 OUNCE PER GALLON	1.25 OUNCES PER GALLON	4 OUNCES PER GALLON
<ul style="list-style-type: none"> • Bordetella bronchispetica (kennel cough) • Salmonella • MRSA • Human Coronavirus (COVID-19) • Avian Influenza • Canine Distemper • Canine Coronavirus • Ringworm 	<ul style="list-style-type: none"> • Canine Adenovirus • Canine parainfluenza virus • Feline Calicivirus • Feline Infectious Peritonitis virus • Feline Panleukopenia virus • Rabies Virus 	<ul style="list-style-type: none"> • Canine Parvovirus • Minute virus of Mice (Parvovirus) • Porcine Parvovirus

*Please refer to Animal Facility Disinfectant product reference sheet for full list of infectious agents.

SURFACE CLEANING

1. Put on safety gloves.
2. Wipe the surface, removing pet hair and soil.
3. Set the spray bottle nozzle to the course or stream setting to minimize migration of the chemical spray.
4. Apply solution by holding spray bottle 6 to 8 inches above the surface.
5. Agitate surface using a disposable wiping towel or brush to remove heavy soil.
6. Allow solution to stand for at least 10-minutes.
7. After 10-minutes use a dry paper towel to remove any excess moisture.
8. Allow the surface to dry before reintroducing animals to the area.

FLOOR CLEANING

1. Put on safety gloves.
2. Place wet floor caution signs to prevent slip-and-falls.
3. Sweep and/or dust to remove all loose debris.
4. Working from the back of the room or area towards the exit, apply a liberal amount of solution onto the floor surface including under counters, cages and along baseboards.
5. Allow the solution to stand for 10 minutes.
6. Allow the floor to dry thoroughly before reintroducing animals to the area.

Rinsing Procedure (if required)

Rinsing is recommended if the floor has a high level of soil, the solution is prepared at 4-ounces per gallon, or if a visible residue builds up over time.

1. Fill a mop bucket with clean water.
2. Using a clean mop head, mop the floor in a side-to-side motion.
3. Dump and replace water when it becomes visibly dirty and rinse mop head.
4. Allow the floor to dry thoroughly before reintroducing animals to the area.



Licensed Veterinary Medical CE Provider – License Number: PVD134
For more information, please call 800.869.4789