

# Animal Facility Disinfectant

Concentrated Disinfectant Cleaner & Deodorizer



Veterinarian type disinfectant, formulated without strong acid or chlorine!

For Use In: Veterinary Hospitals, Kennels, Animal Shelters,  
Animal Life Science Laboratories, Pet Grooming, Retail Pet Shops,  
Equine Hospitals & Stables, Turkey Farms, Swine Quarters and  
Zoos



# Animal Facility Disinfectant

Concentrated Disinfectant Cleaner & Deodorizer

For Use In: Veterinary Hospitals, Kennels, Animal Shelters, Animal Life Science Laboratories, Pet Grooming, Retail Pet Shops, Equine Hospitals & Stables, Turkey Farms, Swine Quarters and Zoos

Veterinarian type disinfectant, formulated without strong acid or chlorine!

This product has been tested and formulated to be efficacious against the following micro-organisms on hard, non-porous surfaces:

#### Bacteria – 1 Ounce Per Gallon

- Pseudomonas aeruginosa*
- Salmonella enterica*
- Staphylococcus aureus*
- Escherichia coli*
- Klebsiella pneumoniae*
- Klebsiella aerogenes*
- Salmonella schottmuelleri*
- Salmonella enterica*
- Shigella dysenteriae*
- Staphylococcus aureus*

#### Bacteria – 4 Ounces Per Gallon

- Bordetella bronchiseptica*

#### Viruses – 1 Ounce Per Gallon

- Influenza A (H1N1) virus*
- Coronavirus 2 (SARS-V) – 1 Minute*
- Bovine Viral Diarrhea*
- Canine Distemper Virus*
- Herpes Simplex Type 1 virus*
- Influenza A2/Japan*
- Norwalk/Norovirus*
- Pseudorabies virus*
- Vaccinia virus*
- Feline Calicivirus*

#### Viruses – 4 Ounces Per Gallon

- Canine Coronavirus*
- Canine Adenovirus*
- Canine Parvovirus*
- Canine Parainfluenza Virus*
- Feline Panleukopenia virus*
- Feline Coronavirus*
- Feline Picornavirus*
- Feline Rhinotracheitis virus*
- Vaccinia virus*

#### Mold and Mildew Control

Use this product at 1 ounce per gallon to control the growth of mold and mildew and their odors on hard, nonporous surfaces. Thoroughly wet all treated surfaces completely. Let air-dry. Repeat application weekly or when growth or odor reappears.

#### Physical Properties

<b>pH of Concentrate</b>	9.5 – 11.5	<b>Flash Point</b>	>200 F
--------------------------	------------	--------------------	--------

#### Animal Care Use Sites:

Veterinary Hospital Waiting	Examination, X-Ray, Operating	Kennels
Life Science Laboratories	Grooming Establishments	Retail Pet Shops
Zoos	Equine Stables/Hospitals	Turkey Farms

#### Animal Care Use Surfaces:

Animal Washing Area	Formica	Plastic (Upholstery)
Cabinets	Furniture	Stainless Steel
Chrome	Glass	Treatment Tables
Desks	Glazed Tile	Telephones/Keypads
Doorknobs & Handles	Granite	Toilet Bowels
Enameled Surfaces	Lift Tables	Urinals
Faucets	Marble	Vinyl (Upholstery)
Feeding Equipment	Watering Equipment	Metal
Washable Walls	Conductive Flooring	Painted Woodwork

#### Poultry Premise Surfaces:

Egg Receiving Area	Tray Dumping Area	Chick Processing Area
Egg Holding Area	Chick Holding Area	Chick Loading Area
Setter Room	Hatchery Room	Poultry Buildings

#### Swine Premise Surfaces:

Farrowing Barns & Areas	Dressing Plants	Blocks
Waters & Feeders	Loading Equipment	Creep Area
Hauling Equipment	Nursery	Chutes

#### \*The AOAC Use Dilution Test Method

The AOAC Use Dilution method is a method of testing the efficacy of disinfectants, originally developed in 1955. Throughout its numerous revisions, it has become the standard for evaluating liquid and dilutable liquid disinfectants for hard surfaces. In particular, this method is specified by the U.S. EPA as the required method for disinfectant claim substantiation.



# Animal Facility Disinfectant

## Concentrated Disinfectant Cleaner & Deodorizer

For Use In: Veterinary Hospitals, Kennels, Animal Shelters, Animal Life Science Laboratories, Pet Grooming, Retail Pet Shops, Equine Hospitals & Stables, Turkey Farms, Swine Quarters and Zoos

**Veterinarian type disinfectant, formulated without strong acid or chlorine!**

### DIRECTIONS FOR USE

**GENERAL CLEANING** – Apply Animal Facility Disinfectant to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low-pressure coarse sprayer. Animal Facility Disinfectant can be diluted at 1 to 4 ounces per gallon (please refer to efficacy list) . Thoroughly wet soiled surface and scrub, as necessary.

**PET ODORS:** Use on carpets, floors, walls, tile, cages, pet beds, litter boxes, mats, floor coverings, area rugs or any surface soiled by a pet. Test a small inconspicuous area first. Blot problem area.

### SITE AND APPLICATION SPECIFIC DIRECTIONS FOR USE

**Veterinary Hospitals – Animal Shelters – Kennels – Grooming Establishments – Retail Pet Shops – Life Science Laboratories – Zoos**

**Disinfection & Virucidal Activity Directions** – For cleaning and disinfecting the following hard nonporous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc., remove all animals and feed from premises. Remove litter, droppings and manure from floors, walls and surfaces occupied or traversed by animals. Saturate surfaces with a use-solution of 4 fluid ounces of Animal Facility Disinfectant per gallon of water and allow the surfaces to remain visibly wet for a period of 10 minutes. Ventilate buildings and closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, automatic feeders, waters and other equipment that dispenses food or water with soap or detergent, and rinse with potable water before reuse.

### **Poultry and Swine Premise Sanitation Directions**

**Site Preparation:** The first step in any on-going sanitation program must be the removal of visible contamination and debris. This may be accomplished using a shovel, broom, or vacuum depending on the area being disinfected. The efficacy of even the most efficient germicidal cleaner is reduced in the presence of visible organic matter. Once the visible debris is eliminated, thoroughly clean all surfaces with enzymatic cleaner and rinse with water.

**Cleaning and Disinfection:** For general cleaning and disinfecting of hard, nonporous surfaces use 4 fluid ounces of Animal Facility Disinfectant per gallon of water. Apply Animal Facility Disinfectant using cloth, mop, hand pump trigger sprayer or low-pressure coarse sprayer to thoroughly wet surface to be disinfected. Allow to remain visibly wet for 10 minutes and then air dry. For visibly soiled areas, a pre-cleaning step is required. Prepare a fresh solution after each use.

### **Farm Premise Disinfection Directions (For Use in Equine, Dairy and Hog Farms)**

Product may only be used on hard, nonporous surfaces for disinfection. Do not use in milking stalls, milking parlors or milk houses.

1. Remove all animals and feed from premises, vehicles and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied by transversed animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent and rinse with water.
5. Saturate all surfaces with a use-solution of 1 to 4 fluid ounces of Animal Facility Disinfectant per gallon of water and allow surface to remain visibly wet for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure.
7. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not hose livestock or employ equipment until treatment has been absorbed, set or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

**Non-Acid Toilet Bowl Disinfection/Cleaner Directions:** Remove visible soil prior to disinfection.

**From concentrate:** Add 1 ounce per gallon to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim, allow to stand for 10 minutes and flush. (For heavy duty use: Add 4 fluid ounces per gallon to the toilet bowl and mix. Brush or swab thoroughly under the rim with a cloth, mop, sponge, or sprayer, allow to stand for 10 minutes and flush.)

**Animal Facility Disinfectant – 32 Oz.  
AcuPro Precision Pour Bottle**  
(Squeeze, dilute and pour)



ITEM NO.:	PACK
V01-32MN	6/32-Ounce

**Animal Facility Disinfectant – 64 Oz.  
eZ Pour Bottle**



ITEM NO.:	PACK
V01-.5MN-005	2/64-Ounce

**ProLoc Wall-Mounted Hospital Station**  
(For use with V01-.5MN Animal Facility Disinfectant)



ITEM NO.:	PACK
V99	Each

**Animal Facility Disinfectant – 64 Oz.  
ProLoc Dispenser Bottle**  
(Fits our ProLoc Dispenser)



ITEM NO.:	PACK
V01-.5MN	4/64-Ounce

**Animal Facility Disinfectant - Gallon  
eZ Pour Bottle**



ITEM NO.:	PACK
V01-14MN	4/1 Gallon

**Animal Facility Disinfectant - 2.5 Gallon  
eZ Pour Bottle**



ITEM NO.:	PACK
V01-25MN	2/2.5 Gallon

**Animal Facility Disinfectant - 30 Gallon  
Drum**



ITEM NO.:	PACK
V01-30MN	30 Gallon

**Animal Facility Disinfectant - 55 Gallon  
Drum**



ITEM NO.:	PACK
V01-55MN	55 Gallon

**Animal Facility Disinfectant - Spray  
Bottle Labels**  
(For Ready-to-Use Animal Facility Disinfectant)



ITEM NO.:	PACK
V01-LABEL	Each

**Animal Facility Disinfectant - Refillable  
Wiping System Canister**  
(With 160 count towel roll)



ITEM NO.:	PACK
PVL-115-012	1 - Canister

**Animal Facility Disinfectant - Refillable  
Wiping System Towel Roll**  
(160 durable wipes)



ITEM NO.:	PACK
PVL-10017R-6	6 – Towel Rolls

**Animal Facility Disinfectant - Refillable  
Wiping System Locking Canister  
Bracket**



ITEM NO.:	PACK
PVL-SS20012	1 – Each

**EPA** Product is tested and approved by the E.P.A.



Product is effective against COVID-19.

